

Vanishing thetanulls and hyperelliptic curves

Olivier Schneider

1st February 2008

Abstract

Let $\mathcal{M}_g(2)$ be the moduli space of curves of genus g with a level-2 structure. We prove that there is always a non hyperelliptic element in the intersection of four thetanull divisors in $\mathcal{M}_6(2)$. We prove also that for all $g \geq 3$, each component of the hyperelliptic locus in $\mathcal{M}_g(2)$ is a connected component of the intersection of $g-2$ thetanull divisors.

1 Introduction

Let C be a curve of genus $g \geq 3$ over $\mathbb{C}$ and $\sigma : \mathbb{Z}^{2g} \longrightarrow H_1(C, \mathbb{Z})$ a symplectic isomorphism. We associate to (C, σ) a period matrix Z in the Siegel upper half space H_g and the numbers

$$\theta[k](Z) := \sum_{r \in \mathbb{Z}^g} \exp \left(\pi \sqrt{-1} \left[{}^t \left(r + \frac{k'}{2} \right) Z \left(r + \frac{k'}{2} \right) + 2 {}^t \left(r + \frac{k'}{2} \right) \frac{k''}{2} \right] \right).$$

for each $k = (k', k'')$ in $(\mathbb{Z}/2\mathbb{Z})^{2g}$ such as $k' \cdot k''$ is even. The vanishing of $\theta[k](Z)$ depends only of $\sigma \pmod{2}$, that is, on the class of (C, σ) in $\mathcal{M}_g(2)$, the moduli space of curves of genus g on $\mathbb{C}$ with a level-2 structure. Thus the zero locus of $\theta[k]$ defines a divisor in $\mathcal{M}_g(2)$ called thetanull divisor of characteristic $[k]$. In genus 3, each thetanull divisor is a component of the hyperelliptic locus in $\mathcal{M}_3(2)$. In genus 4, we know since Riemann that each intersection of two thetanull divisors is an union of hyperelliptic components in $\mathcal{M}_4(2)$. In genus 5, Accola has established in [A1], that with a condition on the three characteristics, the intersection of the corresponding divisors is an union of hyperelliptic components. We propose here to prove that this fails in genus 6 :

Theorem 1.1 *Each sub-variety of $\mathcal{M}_6(2)$ intersection of four thetanull divisors contains an element wich is not hyperelliptic.*

The first step of the proof is to classify the orbits of the action of $\mathrm{Sp}_{2g}(\mathbb{Z}/2\mathbb{Z})$ on the set of quadruplets of thetanull divisors. Afterwards, we finish the proof by verifying that one element of each orbit (and then every quadruplet) defines a subvariety of $\mathcal{M}_6(2)$ which contains a bi-elliptic curve. In view of this result, it is unlikely that hyperelliptic components are intersection of $g - 2$ thetanull divisors in higher genus. Nevertheless, in the last part, we prove the following result :

Theorem 1.2 *Every component of the hyperelliptic locus in $\mathcal{M}_g(2)$ can be defined as a connected component of the intersection of $g - 2$ thetanull divisors.*

2 Preliminaries

In what follows we denote by C a curve of genus g over $\mathbb{C}$. Let $\mathcal{A}_{g,2}$ be the moduli space of g -dimensional principally polarized abelian varieties with a level-2 structure. It can be described as follows (see for instance [BL] chapter 4). Let H_g be the Siegel generalized half-space :

$$H_g = \{Z \in M_g(\mathbb{C}) \mid {}^t Z = Z, \mathrm{Im} Z > 0\}.$$

Then to each Z in H_g corresponds the complex torus $\mathbb{C}^g/(\mathbb{Z}^g + Z\mathbb{Z}^g)$ which comes with a natural principal polarization and a level-2 structure. Moreover we let

$$\Gamma_g(2) := \{M \in \mathrm{Sp}_{2g}(\mathbb{Z}) \mid M \equiv 1_{2g} \pmod{2}\}.$$

this group acts on H_g in the following way :

$$M.Z := (AZ + B)(CZ + D)^{-1}, \quad M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \Gamma_g(2).$$

Then $\mathcal{A}_{g,2}$ is isomorphic to the quotient space $H_g/\Gamma_g(2)$. We denote by $\mathcal{M}_{g,2}$ the space which parametrizes the pairs (C, σ) where C is a curve of genus g and $\sigma : (\mathbb{Z}/2\mathbb{Z})^{2g} \longrightarrow H_1(C, \mathbb{Z}/2\mathbb{Z})$ a symplectic isomorphism. $\mathrm{Sp}_{2g}(\mathbb{Z}/2\mathbb{Z})$ acts on $\mathcal{M}_{g,2}$ in a natural way. Let $k = (k', k'') \in (\mathbb{Z}/2\mathbb{Z})^{2g}$ such as $k'.k''$ is even (in what follows, we will say that k is even). Let

$$\theta[k] : H_g \longrightarrow \mathbb{C}$$

be the function defined by

$$\theta[k](Z) = \sum_{r \in \mathbb{Z}^g} \exp \left(\pi \sqrt{-1} \left[{}^t \left(r + \frac{k'}{2} \right) Z \left(r + \frac{k'}{2} \right) + 2 {}^t \left(r + \frac{k'}{2} \right) \frac{k''}{2} \right] \right).$$

Let M be in $\Gamma_g(2)$. As a consequence of the transformation formula (see for instance [I] page 176), $\theta[k](M.Z)$ is proportionnal to $\theta[k](Z)$. By this fact, the zero locus of $\theta[k]$ defines a divisor of $\mathcal{M}_g(2)$ denoted by $\theta[k]_0$ and called the **thetanull divisor** of characteristic $[k]$. Finally, the natural action of $\mathrm{Sp}_{2g}(\mathbb{Z}/2\mathbb{Z})$ on $(\mathbb{Z}/2\mathbb{Z})^{2g}$ induces an action on these hypersurfaces : For each k in $(\mathbb{Z}/2\mathbb{Z})^{2g}$ wich is even, for each M in $\mathrm{Sp}_{2g}(\mathbb{Z}/2\mathbb{Z})$, we have

$$M.\theta[k]_0 = \theta[M.k]_0.$$

3 Theta characteristics, symplectic torsors

Each concept of this part can be found (for instance) in [S].

Definition 3.1 *Let C be a curve of genus g over $\mathbb{C}$ and K its canonical divisor class. A **theta characteristic** Δ on C is a degree $g-1$ divisor class such as*

$$2\Delta \equiv K$$

*Moreover Δ is called **even** (resp **odd**) if $h^0(\Delta) \equiv 0 \pmod{2}$ (resp $1 \pmod{2}$).*

Let $\mathcal{S}(C)$ be the set of theta characteristics on C and let $\mathcal{S}^+(C)$ be the set of even ones.

Definition 3.2 *Let $(J, .)$ be a symplectic pair (that is a $\mathbb{Z}/2\mathbb{Z}$ -vector space endowed with a non degenerate, alternate, bilinear form), we say that a pair (S, Q) is a symplectic torsor over $(J, .)$ if there is simply transitive action of J on S denoted $+$ and a mapping $Q : S \longrightarrow \mathbb{Z}/2\mathbb{Z}$ having the property $\forall s \in S, \forall j_1 \in J, \forall j_2 \in J,$*

$$Q(s) + Q(j_1 + s) + Q(j_2 + s) + Q(j_1 + j_2 + s) = j_1.j_2 \quad (*)$$

Example 3.3 Let W be a set of order $2g+2$. We denote by $P_2(W)$ the set of partitions on W into two subsets. For each subset A of W , we denote by $\bar{A}$ the element $(A; \mathbb{C}_W A)$ of $P_2(W)$. One has

1. An addition on $P_2(W)$: For each $\bar{A}$ and $\bar{B}$ in $P_2(W)$

$$\bar{A} + \bar{B} := \overline{A \cup B - A \cap B}$$

2. A map $p : P_2(W) \longrightarrow \mathbb{Z}/2\mathbb{Z}$ defined by

$$p(\bar{A}) = |A| \pmod{2}.$$

We denote by $P_2^+(W)$ (resp $P_2^-(W)$) the set $p^{-1}(0)$ (resp $p^{-1}(1)$).

3. A map $e : P_2^+(W) \times P_2^+(W) \longrightarrow \mathbb{Z}/2\mathbb{Z}$ defined by

$$e(\bar{A}; \bar{B}) := |A \bigcup B| \pmod{2}.$$

If g is even (resp odd) then we have a quadratic form defined on $P_2^-(W)$ (resp on $P_2^+(W)$) by

$$Q_-(\bar{A}) = \frac{|A| + 1}{2} \pmod{2}$$

$$\left(\text{resp } Q_+(\bar{A}) = \frac{|A|}{2} \pmod{2} \right).$$

$(P_2^-(W), Q_-)$ (resp $(P_2^+(W), Q_+)$) is a symplectic torsor over $(P_2^+(W), e)$ denoted by $T(W)$.

Example 3.4 For each symplectic pair $(J, \cdot)$, the set of quadratic forms

$$Quad(J) := \{q : J \longrightarrow \mathbb{Z}/2\mathbb{Z} \mid \forall j_1 \in J, \forall j_2 \in J, q(j_1) + q(j_2) + q(j_1 + j_2) = j_1 \cdot j_2\}$$

endowed with the mapping $\text{Arf} : Quad(J) \longrightarrow \mathbb{Z}/2\mathbb{Z}$ (see for instance [Sc]) is a symplectic torsor over $(J, \cdot)$. Moreover we have the following property : $\forall q \in Quad(J), \forall j \in J,$

$$\text{Arf}(j + q) = \text{Arf}(q) + q(j)$$

Proposition 3.5 *Let $(\mathcal{J}_2(C), \cdot)$ be the $\mathbb{Z}/2\mathbb{Z}$ -vector space of order two points of the Jacobian of C endowed with the intersection pairing. If we denote by Q the mapping*

$$\begin{aligned} Q : \mathcal{S}(C) &\longrightarrow \mathbb{Z}/2\mathbb{Z} \\ \Delta &\longmapsto h^0(\Delta) \pmod{2} \end{aligned}$$

then $(\mathcal{S}(C), Q)$ is a symplectic torsor over $(\mathcal{J}_2(C), \cdot)$.

Remark 3.6 Let C be an hyperelliptic curve. If we denote by $W(C)$ the set of its Weierstrass points, then we have a canonical isomorphism of symplectic torsors

$$\widetilde{\beta}_C : T(W(C)) \longrightarrow (\mathcal{S}(C), Q)$$

compatible with the canonical symplectic isomorphism

$$\beta_C : (P_2^+(W(C)), e) \longrightarrow (\mathcal{J}_2(C), \cdot).$$

Finally, we recall the result established by Mumford in [M] :

Proposition 3.7 *The map*

$$\begin{array}{ccc} (\mathcal{S}(C), Q) & \longrightarrow & (Quad(\mathcal{J}_2(C)), \text{Arf}) \\ \Delta & \longmapsto & \left(\begin{array}{ccc} q_\Delta : \mathcal{J}_2(C) & \longrightarrow & \mathbb{Z}/2\mathbb{Z} \\ j & \longmapsto & Q(\Delta) + Q(j + \Delta) \end{array} \right) \end{array}$$

is an isomorphism of symplectic torsors. In particular it preserves the parity.

4 Proof of theorem 1.1

4.1 Orbits of the action of $\text{Sp}_{12}(\mathbb{Z}/2\mathbb{Z})$ on the sets of four distinct even theta characteristics

Let C be a curve of genus 6 over $\mathbb{C}$. For each set $\{\Delta_i\}_{i=1\dots 4}$ of four distinct even theta characteristics on C , we can associate the set $\{\delta_i\}_{i=1\dots 4}$ of theta characteristics on C , defined as

$$\delta_i \equiv 2K - \sum_{j \neq i} \Delta_j.$$

Proposition 4.1 *Let C be a curve of genus 6 over $\mathbb{C}$. The action of $\text{Sp}_{12}(\mathbb{Z}/2\mathbb{Z})$ on the sets of four even theta characteristics has four orbits $A_1, \dots, A_4$, given respectively by the following conditions :*

1. $\sum_{i=1}^4 \Delta_i \equiv 2K$.
2. $\{\Delta_i\}_{i=1\dots 4}$ is not in A_1 and all the δ_i 's are even.
3. Exactly two of the δ_i 's are even.
4. All the δ_i 's are odd.

By Proposition 3.7, we can study the action of $\text{Sp}_{12}(\mathbb{Z}/2\mathbb{Z})$ on the sets of four distinct even quadratic forms on $\mathcal{J}_2(C)$:

Lemma 4.2 *Let C be a curve of genus 6 over $\mathbb{C}$. Let $\{q, q+a_1, q+a_2, q+a_3\}$ and $\{q, q+a'_1, q+a'_2, q+a'_3\}$ be two sets of four distinct even quadratic forms on $\mathcal{J}_2(C)$ so that*

$$\dim \langle a_1, a_2, a_3 \rangle = \dim \langle a'_1, a'_2, a'_3 \rangle .$$

If it exists $\sigma \in \mathfrak{S}_3$ so that

$$\forall k, l \in \{1, 2, 3\}, \quad a_k \cdot a_l = a'_{\sigma(k)} \cdot a'_{\sigma(l)},$$

then these two sets are in the same orbit of the action of $\mathrm{Sp}_{12}(\mathbb{Z}/2\mathbb{Z})$ on the sets of four distinct even quadratic forms on $\mathcal{J}_2(C)$.

Proof : By taking into account the parity conditions on the quadratic forms, we have :

$$\begin{aligned} q(a_1) &= q(a'_1) \\ q(a_2) &= q(a'_2) \\ q(a_3) &= q(a'_3). \end{aligned}$$

Moreover, $\dim \langle a_1, a_2, a_3 \rangle = \dim \langle a'_1, a'_2, a'_3 \rangle$ and it exists $\sigma \in \mathfrak{S}_3$ so that

$$\forall k, l \in \{1, 2, 3\}, \quad a_k \cdot a_l = a'_{\sigma(k)} \cdot a'_{\sigma(l)}.$$

By Witt's theorem (see for instance [Sc]) these conditions are equivalent to :

$$\exists f \in \mathrm{O}(q) \text{ so that } \forall k \in \{1, 2, 3\}, \quad f(a_k) = a'_{\sigma(k)}.$$

Finally, this is equivalent to the existence of f in $\mathrm{O}(q)$ so that

$$\begin{aligned} q_1 &= q'_{i_1} \circ f^{-1} \\ q_2 &= q'_{i_2} \circ f^{-1} \\ q_3 &= q'_{i_3} \circ f^{-1}. \quad \square \end{aligned}$$

Proof of Proposition 4.1 : As the action preserves the parity, each orbit is contained in one of the A_i 's. By this fact, we have to show the transitivity of this action on these sets :

Let $\Delta_1, \Delta_2, \Delta_3$ and Δ_4 be four even theta characteristics on C . Let q_1, q_2, q_3 and q_4 be the quadratic forms associated to these theta characteristics. For each i in $\{1, 2, 3\}$, let a_i be the element of $\mathcal{J}_2(C)$ so that

$$q_i = q_4 + a_i.$$

If a_1, a_2 and a_3 are linearly dependent then $\Delta_4 + \Delta_1 \equiv \Delta_2 + \Delta_3$.

If a_1, a_2 and a_3 are linearly independent then we have :

$$\forall \{i, j\} \subseteq \{1, 2, 3\},$$

$$a_i \cdot a_j = 0 \Leftrightarrow \Delta_4 + \Delta_i - \Delta_j \text{ even.}$$

Thus, we have :

$$\begin{aligned}
\{\Delta_i\}_{i=1\dots 4} \in A_1 &\Leftrightarrow a_1, a_2 \text{ and } a_3 \text{ linearly dependent.} \\
\{\Delta_i\}_{i=1\dots 4} \in A_2 &\Leftrightarrow \dim \langle a_1, a_2, a_3 \rangle = 3 \text{ and } \forall i \in \{1, 2, 3\}, a_i \cdot a_j = 0 \\
\{\Delta_i\}_{i=1\dots 4} \in A_3 &\Leftrightarrow \dim \langle a_1, a_2, a_3 \rangle = 3 \text{ and} \\
&\quad \text{Card}(\{\{i, j\} \subset \{1, 2, 3\} \mid a_i \cdot a_j = 1\}) = 1 \text{ or } 2 \\
\{\Delta_i\}_{i=1\dots 4} \in A_4 &\Leftrightarrow \dim \langle a_1, a_2, a_3 \rangle = 3 \text{ and } \forall i \in \{1, 2, 3\}, a_i \cdot a_j = 1
\end{aligned}$$

Let q_1^*, q_2^*, q_3^* and q_4^* be four even quadratic forms on $\mathcal{J}_2(C)$ corresponding to $\{\Delta_i^*\}_{i=1\dots 4}$, an other set of even theta characteristics on C . The transitivity of the action (on even quadratic forms) implies that there exists M in $\text{Sp}_{12}(\mathbb{Z}/2\mathbb{Z})$, so that $M \cdot q_4^* = q_4$. If we put

$$\begin{aligned}
q_1' &:= M \cdot q_1^* \\
q_2' &:= M \cdot q_2^* \\
q_3' &:= M \cdot q_3^*,
\end{aligned}$$

then $\{q_1', q_2', q_3', q_4\}$ is in the same orbit as $\{q_1^*, q_2^*, q_3^*, q_4^*\}$.

By this fact, if $\{\Delta_i\}_{i=1\dots 4}$ and $\{\Delta_i^*\}_{i=1\dots 4}$ are both in A_1, A_2, A_4 , then the transitivity is obvious by lemma 4.2.

Now let us suppose that $\{\Delta_i\}_{i=1\dots 4}$ and $\{\Delta_i^*\}_{i=1\dots 4}$ are in A_3 . For each i in $\{1, 2, 3\}$, let a_i' be the element of $\mathcal{J}_2(C)$ so that

$$q_i' = q_4 + a_i'.$$

If $\text{Card}(\{\{i, j\} \subset \{1, 2, 3\} \mid a_i \cdot a_j\}) = \text{Card}(\{\{i, j\} \subset \{1, 2, 3\} \mid a_i' \cdot a_j'\})$, then we can conclude again with lemma 4.2. If not, we can suppose for instance, that

$$\begin{aligned}
a_1 \cdot a_2 &= 1 & a_1' \cdot a_2' &= 1 \\
a_1 \cdot a_3 &= 0 & \text{and } a_1' \cdot a_3' &= 1 \\
a_2 \cdot a_3 &= 0 & a_2' \cdot a_3' &= 0.
\end{aligned}$$

By lemma 4.2, there exist a_2'' and a_3'' in $\mathcal{J}_2(C)$ so that

$$\begin{aligned}
a_2 \cdot a_2'' &= 1 \\
a_2 \cdot a_3'' &= 1 \\
a_2'' \cdot a_3'' &= 0,
\end{aligned}$$

and $\{q_4 + a_2, q_4 + a_2'', q_4 + a_3'', q_4\}$ is in the same orbit as $\{q_4 + a_1', q_4 + a_2', q_4 + a_3', q_4\} = \{q_1', q_2', q_3', q_4\}$. If we put

$$q := q_4 + a_2,$$

then

$$\begin{aligned}\{q_1, q_2, q_3, q_4\} &= \{q + (a_1 + a_2), q, q + (a_3 + a_2), q + a_2\} \\ \{q_4 + a_2, q_4 + a_2'', q_4 + a_3'', q_4\} &= \{q, q + (a_2 + a_2''), q + (a_2 + a_3''), q + a_2\}.\end{aligned}$$

We can verify that

$$\begin{aligned}(a_1 + a_2).(a_2 + a_3) &= 1 & a_2.(a_2 + a_2'') &= 1 \\ (a_1 + a_2).a_2 &= 1 & \text{and} & a_2.(a_2 + a_3'') &= 1 \\ (a_2 + a_3).a_2 &= 0 & (a_2 + a_2'').(a_2 + a_3'') &= 0.\end{aligned}$$

and conclude with the lemma 4.2. $\square$

4.2 Theta characteristics on bi-elliptic curves

Let C be a bi-elliptic curve of genus 6. By definition this means that there exists an elliptic curve X and a degree 2 morphism :

$$\pi : C \longrightarrow X$$

The Riemann-Hurwitz theorem implies that π has 10 ramification points denoted $R_1, \dots, R_{10}$.

Lemma 4.3 ([A2]) *A general bi-elliptic curve has forty even, effective theta characteristics of the form*

$$R_i + \pi^*(D_i)$$

where R_i is one of the ramification points of π and D_i is a degree two divisor on X .

Remark 4.4 • R_i is the fixed point of the linear system $|R_i + \pi^*(D_i)|$.

- $\mathcal{J}(X)$ has 3 non zero points of order 2 denoted by F_1, F_2 and F_3 . If for each i in $\{1, \dots, 10\}$, we choose a divisor D_i so that $R_i + \pi^*D_i$ is an even theta characteristic on C , then the three other even, effective theta characteristics with fixed point R_i will be

$$R_i + \pi^*(D_i + F_j) \quad j = 1 \dots 3$$

Lemma 4.5 *Let $\Delta_1 = R_1 + \pi^*(D_1)$, $\Delta_2 = R_2 + \pi^*(D_2)$ and $\Delta_3 = R_3 + \pi^*(D_3)$ be three even effective theta characteristics on a general bi-elliptic curve C . One has*

$$\Delta_1 + \Delta_2 - \Delta_3 \text{ even} \Leftrightarrow \begin{cases} \text{two at least of} \\ \text{the } R_i \text{ are the same} \end{cases}$$

Proof : First of all, let us notice that for each j, k, l so that $\{j, k, l\} = \{1, 2, 3\}$,

$$\Delta_1 + \Delta_2 - \Delta_3 = R_j + R_k - R_l + \pi^* D_j + \pi^* D_k - \pi^* D_l$$

then $D_j + D_k - D_l \equiv E_l$ where E_l is a degree two, effective divisor on X ; thus $(\Leftarrow)$ is evident.

Now, if we call P_l , the point on X so that $\pi^* P_l = 2R_l$, as on an elliptic curve every linear system of degree d is of dimension d , there exists Q_l , a point on X so that $P_l + Q_l \equiv E_l$; thus

$$\begin{aligned} R_j + R_k - R_l + \pi^* D_j + \pi^* D_k - \pi^* D_l &\equiv R_j + R_k - R_l + \pi^* E_l \\ &\equiv R_j + R_k - R_l + \pi^* P_l + \pi^* Q_l \\ &\equiv R_1 + R_2 + R_3 + \pi^* Q_l \end{aligned}$$

But then, by lemma 4.5, for a general bi-elliptic curve, if we are considering, for instance, that R_1 is the fixed point of the even theta characteristic $\Delta_1 + \Delta_2 - \Delta_3$, then $R_2 + R_3 + \pi^* Q_l$ must be the pullback of an effective degree two divisor. This is possible if and only if $R_2 = R_3$. $\square$

With this lemma, one checks easily the next result which ends our proof.

Proposition 4.6 *Let C be a smooth curve of genus 6 with a degree two morphism*

$$\pi : C \longrightarrow X$$

onto an elliptic curve X . Let $R_1, \dots, R_{10}$ be the ramification points of this morphism and let $D_1, \dots, D_{10}$ be some degree two effective divisors on X so that

$$\{R_i + \pi^* D_i \mid i \in \{1, \dots, 10\}\}$$

is a set of the even, effective theta characteristic on C . The following sets are in the four orbits of the action of $\mathrm{Sp}_{12}(\mathbb{Z}/2\mathbb{Z})$ on the sets of four distinct even theta characteristics on C :

$$\begin{aligned} \{R_1 + \pi^* D_1, R_2 + \pi^*(D_2 + F_1), R_1 + \pi^*(D_1 + F_1), R_2 + \pi^* D_2\} &\in A_1 \\ \{R_1 + \pi^* D_1, R_2 + \pi^*(D_2 + F_2), R_1 + \pi^*(D_1 + F_1), R_2 + \pi^* D_2\} &\in A_2 \\ \{R_1 + \pi^* D_1, R_2 + \pi^*(D_2), R_3 + \pi^*(D_3), R_1 + \pi^*(D_2 + F_1)\} &\in A_3 \\ \{R_1 + \pi^* D_1, R_2 + \pi^* D_2, R_3 + \pi^* D_3, R_4 + \pi^* D_4\} &\in A_4. \end{aligned}$$

5 Proof of theorem 1.2

First of all, we need to recall a result given by Teixidor I Bigas in [T] :

Let $\mathfrak{S}_1$ be the scheme wich parametrizes the pairs (C, Δ) , where C is curve of genus g on $\mathbb{C}$ and Δ is a theta characteristic of projective dimension 1 on C . For each $t = (C, \Delta)$ in $\mathfrak{S}_1$, there is an injective map :

$$f : T_{\mathfrak{S}_1(t)} \longrightarrow H^1(T_C)$$

As by Serre duality $H^1(T_C)$ is dual to $H^0(2K)$, we have :

Lemma 5.1 (*Teixidor I Bigas*) *If F is the fixed part of the linear system $|\Delta|$ and R the ramification divisor of the corresponding morphism $\varphi_\Delta : C \longrightarrow \mathbb{P}^1$ then $\text{Im}(f) = (\omega)^\perp$, where ω is an element of $H^0(2K)$ with divisor $R + 2F$.*

By using the particular expression of the theta characteristics on an hyperelliptic curve (see [A,C,G,H] p 288), this lemma has the following consequence : Let C be an hyperelliptic curve of genus $g \geq 3$, $W := \{p_1, \dots, p_{2g+2}\}$ the set of its Weierstrass points, H its hyperelliptic divisor, then for each even theta characteristic Δ on C of dimension 1,

$$\Delta = H + p_{i_1} + \dots + p_{i_{g-3}}.$$

$\text{Im}(f)$ is the orthogonal of ω so that

$$\text{div}(\omega) = p_1 + \dots + p_{2g+2} + 2(p_{i_1} + \dots + p_{i_{g-3}})$$

Proposition 5.2 *Let (C, σ) be an hyperelliptic element of $\mathcal{M}_g(2)$ ($g \geq 3$). Let H be the hyperelliptic divisor on C . We choose $p_1, \dots, p_{g-2}$, $g-2$ Weierstrass points on C and let E be the divisor $p_1 + \dots + p_{g-2}$. Then the $g-2$ thetanull divisors associated to the theta characteristics $\{H + E - p_1, \dots, H + E - p_{g-2}\}$ intersect transversally at (C, σ) .*

Proof : By the last remark, we have to prove that the linear subsystem of $H^0(2K)$ generated by

$$\{F_k = p_1 + \dots + p_{2g+2} + 2(E - p_k) \mid k = 1, \dots, g-2\}$$

has rank $g-2$. Let us prove that for each k in $\{2, \dots, g-2\}$, F_k is not in the linear system generated by $F_1, \dots, F_{k-1}$. Let C be the Riemann surface

$$y^2 = \prod_{i=1}^{2g+2} (x - x_i) \quad , \quad x : C \longrightarrow \mathbb{P}^1$$

so that for each i in $\{1, \dots, 2g+2\}$, $x(p_i) = x_i$. We have then to verify that there does not exist $(\lambda_1 : \dots : \lambda_{k-1})$ in $\mathbb{P}^{k-2}$ so that $2(E - p_k)$ is the divisor f^*0 where

$$f = \lambda_1(x-x_2) \dots (x-x_{g-2}) + \dots + \lambda_{k-1}(x-x_1) \dots (x-x_{k-2})(x-x_k) \dots (x-x_{g-2})$$

Let us suppose it is false. As $2(E - p_k)$ is $2p_1 + \dots + 2p_{k-1} + 2p_{k+1} + \dots + 2p_{g-2}$ this implies

$$f(x_1) = \lambda_1(x_1 - x_2)(x_1 - x_2) \dots (x_1 - x_{g-2}) = 0 \Rightarrow \lambda_1 = 0$$

then

$$f(x_2) = 0 \Rightarrow \lambda_2 = 0$$

then

$$\vdots$$

$$\begin{aligned} f(x_{k-1}) &= \lambda_{k-1}(x_{k-1} - x_1) \dots (x_{k-1} - x_{k-2})(x_{k-1} - x_k) \dots (x_{k-1} - x_{g-2}) = 0 \\ &\Rightarrow \lambda_{k-1} = 0 \end{aligned}$$

Finally we would have $\lambda_1 = \dots = \lambda_{k-1} = 0$ which it is absurd. $\square$

Each connected component Y of $\mathcal{H}_g(2)$ is uniquely determined by a symplectic isomorphism

$$c_Y : (\mathbb{Z}/2\mathbb{Z})^{2g} \longrightarrow P_2^+(\{1, \dots, 2g+2\})$$

and an isomorphism between symplectic torsors

$$\widetilde{c}_Y : (\mathbb{Z}/2\mathbb{Z})^{2g} \longrightarrow T(\{1, \dots, 2g+2\})$$

in the following way :

Let U be the open subvariety of $\mathbb{C}^{2g+2}$ consisting of points with distinct coordinates. To each point $\xi = (\xi_1, \dots, \xi_{2g+2})$ in U , we associate the hyperelliptic curve $C(\xi)$ with Weierstrass points $\xi_1, \dots, \xi_{2g+2}$. We denote by $W(\xi)$ this set. For each ξ in U , the bijection $[1, \dots, 2g+2] \longrightarrow [\xi_1, \dots, \xi_{2g+2}]$ induces a symplectic isomorphism

$$\alpha_\xi : P_2^+(\{1, \dots, 2g+2\}) \longrightarrow P_2^+(W(\xi))$$

and an isomorphism between symplectic torsors :

$$\widetilde{\alpha}_\xi : T(\{1, \dots, 2g+2\}) \longrightarrow T(W(\xi)).$$

Y is the subspace of $\mathcal{M}_g(2)$ defined as

$$\{(C(\xi), \sigma_\xi) \in \mathcal{H}_{g,2} \mid \xi \in U\},$$

so that for each ξ in U , the following diagrams

$$\begin{array}{ccccc} P_2^+(\{1, \dots, 2g+2\}) & \xrightarrow{\alpha_\xi} & P_2^+(C(\xi)) & \xrightarrow{\beta_{C(\xi)}} & \mathcal{J}_2(C(\xi)) \\ & \nwarrow c_Y & & \nearrow \sigma_\xi & \\ & & (\mathbb{Z}/2\mathbb{Z})^{2g} & & \end{array}$$

$$\begin{array}{ccccc} T(\{1, \dots, 2g+2\}) & \xrightarrow{\widetilde{\alpha}_\xi} & T(W(\xi)) & \xrightarrow{\widetilde{\beta_{C(\xi)}}} & \mathcal{S}(C(\xi)) \\ & \nwarrow \widetilde{c}_Y & & \nearrow \widetilde{\sigma}_\xi & \\ & & (\mathbb{Z}/2\mathbb{Z})^{2g} & & \end{array}$$

are commutative. Now let ξ_0 be in U . Let us choose $g-2$ thetanull divisors which correspond on $C(\xi_0)$, to $g-2$ theta characteristics in the configuration of Proposition 5.2. By the last diagram, one sees that for each ξ in U , these $g-2$ thetanull divisors correspond to $g-2$ theta characteristics on $C(\xi)$ in the same configuration as on $C(\xi_0)$. By this fact, Y is a component of the intersection of these $g-2$ thetanull divisors. $\square$

Acknowledgment

I would like to thank my thesis advisor A. Beauville for making useful suggestions that lead to the results of this paper.

References

- [A1] R.ACCOLA
Some loci in Teichmüller space for genus five defined by vanishing theta nulls . Contributions to analysis (a collection of papers dedicated to Lipman Bers), pp.11–18. Academic Press, New York, 1974.

- [A2] R.ACCOLA
Some loci in Teichmüller space for genus six defined by vanishing thetanulls . Differential geometry and complex analysis, 49–63, Springer, Berlin, 1985.

- [A,C,G,H] E.ARBARELLO , M.CORNALBA , P.GRIFFITHS , J.HARRIS
Geometry of algebraic curves. Vol. I. Grundlehren der Mathematischen Wissenschaften, 267. Springer-Verlag, New York, 1985.
- [BL] C.BIRKENHAKE , H.LANGE
Complex abelian varieties. Grundlehren der Mathematischen Wissenschaften, 302. Springer-Verlag, Berlin, 1992.
- [D] R.DONAGI
Big Schottky. Invent. Math. 89 (1987), no. 3, 569–599.
- [I] J.IGUSA
Theta functions. Die Grundlehren der mathematischen Wissenschaften, 194. Springer-Verlag, New York-Heidelberg, 1972.
- [M] D.MUMFORD
Theta characteristics of an algebraic curve. Ann. Sci. Ecole Norm. Sup. (4) 4 (1971), 181–192.
- [S] N.SAAVEDRA RIVANO
Finite geometries in the theory of theta characteristics. Enseignement Math. (2) 22 (1976), no. 3-4, 191–218.
- [Sc] W.SCHARLAU
Quadratic and Hermitian forms. Grundlehren der Mathematischen Wissenschaften, 270. Springer-Verlag, Berlin, 1985.
- [Ts] S.TSUYUMINE
Thetanullwerte on a moduli space of curves and hyperelliptic loci. Math. Z. 207 (1991), no. 4, 539–568.
- [T] M.TEIXIDOR I BIGAS
Half-canonical series on algebraic curves. Trans. Amer. Math. Soc. 302 (1987), no. 1, 99–115.